



**Australian Government**  
**Australian Pesticides and  
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL PRODUCT  
FOR CONTROL OF ANTHRACNOSE IN ALBUS LUPINS**

**PERMIT NUMBER – PER82226**

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

**THIS PERMIT IS IN FORCE FROM 21 SEPTEMBER 2016 TO 31 OCTOBER 2029**

**Permit Holder:**

GRAIN PRODUCERS AUSTRALIA LIMITED  
12 Cooma Road  
BRAIDWOOD NSW 2622

**Persons who can use the product under this permit:**

Persons generally.

## CONDITIONS OF USE

### Product to be used:

AMISTAR 250 SC FUNGICIDE (APVMA No. 58340)

PLUS OTHER REGISTERED PRODUCTS

Containing: 250 g/L AZOXYSTROBIN as the only active constituent

### RESTRAINTS:

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

#### *Spray drift restraints:*

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

#### **Buffer zones for boom sprayers**

Application rate	Boom height over the canopy	Mandatory downwind buffer zones (metres)
		Natural aquatic areas
500 mL/ha	0.5 m or lower	0
	1.0 m or lower	20

DO NOT apply by aircraft unless the following requirements are met:

- spray droplets not smaller than a MEDIUM (fixed-wing application) or (helicopter application) spray droplet size category
- for maximum release heights above the target canopy of 3m or 25% of wingspan or 25% of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

#### **Buffer zones for aircraft**

Type of aircraft	Mandatory downwind buffer zones (metres)
	Natural aquatic areas
Fixed wing	90
Helicopter	0

**Directions for Use:**

Crop Type	Target Disease	Application Rate
ALBUS LUPINS	Anthrachnose ( <i>Colletotrichum lupini</i> )	500 mL product / ha

**Critical Use Comments:**

- Apply up to two (2) foliar applications per season. The first application must be applied before disease is established and no later than the onset of flowering. A second application can be made 10-14 days after the first application.
- Add a non-ionic surfactant at the recommended label rate.
- Apply using calibrated boom sprayer or aerial equipment.
- Refer to product label for information on resistance management.
- Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

**Ground Application:**

- Apply as a foliar spray in a spray volume of 50-100 L/ha.

**Aerial Application:**

- Product must only be applied with aircraft fitted with accurately calibrated equipment. Apply a minimum total spray volume of 30 L/ha with nozzles (e.g. Micro® rotary atomisers, CP nozzles or conventional hydraulic nozzles) set to a minimum spray quality of MEDIUM according to ASAE S572 definition for standard nozzles.

**Safety directions:**

Will irritate the eyes. Avoid contact with eyes. If product in eyes, wash it out immediately with water. When opening the container and preparing spray wear cotton overalls, buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves and goggles or safety glasses. Wash hands after use. After each day's use, wash gloves and goggles or safety glasses and contaminated clothing.

**Withholding Periods:**

Harvest: DO NOT harvest for 4 WEEKS after last application.

Grazing: DO NOT graze or cut for livestock feed for 4 WEEKS after last application.

**Jurisdiction:**

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their *Control-of-Use* legislation means a permit is not required to legalise this off-label use in that State.

**Additional Conditions:**

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the *Agricultural and Veterinary Chemicals Code Act 1994*, in accordance with the obligation imposed by that section.

### ***Export of treated produce***

Maximum Residue Limits (MRLs) have been established to allow treated produce to be used for human or animal consumption. MRLs can be found in the *Agricultural and Veterinary Chemicals (MRL Standard for Residues of Chemical Products) Instrument 2023*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, due account should be taken of the residue limits/tolerances of importing countries. Residues must not exceed requirements of the importing country.

Issued by Australian Pesticides and Veterinary Medicines Authority

Note: 02/07/2019. Expiry date extended to 31/10/2024. Non-technical edits made to 'Critical Use Comments' and 'Export of treated produce' statements. Issued as version 2.

Note: 30/09/2024 - Permit updated to include Restraints. Spray drift restraints and buffer zones updated. WHPs stated as 'WEEKS' rather than 'DAYS'. Additional critical use comment added. Safety directions added. Additional conditions updated to current standards. S161 statement added. Permit expiry extended to 31/10/2029. Permit issued as Version 3.