



**PERMIT TO ALLOW MINOR USE OF AN AGVET CHEMICAL  
PRODUCT FOR THE CONTROL OF FUNGAL DISEASES IN SILVERBEET,  
SPINACH, CHICORY, ENDIVE, PARSLEY and CELERIAC and FOR THE  
CONTROL OF RING SPOT IN BRASSICA LEAFY VEGETABLES.**

**PERMIT NUMBER –PER87973**

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

**THIS PERMIT IS IN FORCE FROM 27 AUGUST 2020 TO 31 AUGUST 2025.**

**Permit Holder:**

HORTICULTURE INNOVATION AUST LTD  
Level 7, 141 Walker Street  
NORTH SYDNEY NSW 2060.

**Persons who can use the product under this permit:**

Persons generally.

## CONDITIONS OF USE

### Product to be used:

SCORE FOLIAR FUNGICIDE (APVMA No. 45810).

PLUS OTHER REGISTERED PRODUCTS

Containing: 250 g/L DIFENOCONAZOLE as the only active constituent.

IMTRADE DEFEND 400 EC FUNGICIDE (APVMA No. 63825).

PLUS OTHER REGISTERED PRODUCTS

Containing: 400 g/L DIFENOCONAZOLE as the only active constituent.

### Restrains:

In protected cropping situations, additional precautions are to be complied with to minimise exposure of all workers, including works not involved in handling the products:

- Whenever possible, evacuate greenhouse before spraying, and do not allow re-entry before spray has settled.
- In larger greenhouses, if complete evacuation of the structure is not possible: DO NOT spray within 20 metres of other personnel.

### Directions for Use:

Crop	Pest	Rate	Critical Comments
Brassica Leafy Vegetables (field and protected)	Ring spot, Alternaria leaf spot ( <i>Mychosphaerella brassicicola</i> )	<u>250g/L product</u> 400 mL/ha  <u>400 g/L products</u> 250 mL/ha	Use sufficient water to obtain thorough and uniform coverage of foliage. DO NOT apply more than three (3) applications per crop at a minimum re-treatment interval of 14-21 days. Apply at first sign of disease by foliar ground boom spray or equivalent only. DO NOT apply more than two (2) consecutive sprays of a difenoconazole based product. Alternate with a recommended protectant fungicide from a different chemical MoA group for at least one application before continuing with a difenoconazole product alone, or in a tank mix.
Celeriac (field grown)	Cercospora early blight (leaf spot) ( <i>Cercospora apii</i> ) and Septoria late blight (leaf blight) ( <i>Septoria apiicola</i> )	<u>250 g/L products</u> 300 mL/ha  <u>400 g/L products</u> 188 mL/ha	Use in a protectant program prior to diseases occurring. DO NOT apply more than six (6) applications per crop at a minimum retreatment interval of 7 days. DO NOT apply in volumes less than 100L/ha. Apply by foliar ground boom sprayer or equivalent only. DO NOT apply more than two (2) consecutive sprays of a difenoconazole based product. Alternate with a recommended

			protectant fungicide from a different chemical MoA group for at least one application before continuing with a difenoconazole product alone, or in a tank mix.
Chicory, Endive, Silverbeet and Spinach (field and protected)	Powdery Mildew ( <i>Erysiphe betae</i> )	<p><u>250g/L products</u> 50 mL/100 L when spray volume exceeds 800 L/ha or 500 mL/ha when spray volume is less than 800 L.</p> <p><u>400 g/L products</u> 31.25 mL/100 L for when spray volume exceeds 800 L/ha or 313 mL/ha when spray volume is less than 800 L.</p>	Apply at first sign of disease by foliar application via boom spray or equivalent only. DO NOT apply more than two (2) applications per crop with a minimum re-treatment interval of 10 days. DO NOT apply in volumes less than 500 L/ha.
Parsley (field and protected)	Alternaria leaf blight ( <i>Alternaria petroselini</i> ) and Cerospora leaf spot ( <i>Cercospora</i> spp.)	<p><u>250 g/L products</u> 300 mL/ha</p> <p><u>400 g/L products</u> 188 mL/ha</p>	Use in a protectant program before diseases occur. Apply at 7 day intervals when conditions favour disease development by foliar application via boom spray or equivalent only. Use sufficient water to obtain thorough and uniform coverage. DO NOT apply more than three (3) applications of difenoconazole per crop with a minimum re-treatment interval of 7 days. DO NOT apply more than two (2) consecutive sprays of difenoconazole alone (then apply a recommended non-group C fungicide or tank-mix difenoconazole with a recommended protectant fungicide for at least one application).

**Withholding periods:**

Brassica leafy vegetables

Harvest: DO NOT harvest for 3 days after application.

Chicory, celeriac, endive, silverbeet, spinach and parsley

Harvest: DO NOT harvest for 7 days after application.

Grazing: DO NOT graze or cut treated area for stock food.

**Jurisdiction:**

All States and Territories, except VIC.

(Note: Victoria is not included in this permit because their ‘control-of-use’ legislation means that a permit is not required to legalise this off-label use in Victoria).

**Additional Conditions:**

This Permit provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit.

*To Avoid Crop Damage*

The sensitivity of some crops and varieties to be treated under this permit has not been fully evaluated under all growing conditions. It is advisable, therefore, to only treat a small area of crop to ascertain their reaction before treating a larger areas.

*Protection of Wildlife, Fish, Crustaceans and Environment*

**Dangerous to aquatic organisms.**

Do not spray within 5 metres downwind of water bodies.

Do not apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, water bodies or watercourses.

Do not contaminate dams, drains, streams, rivers or waterways with the chemical or used containers.

*Export of treated produce.*

Temporary Maximum Residue Limits (TMRLs) have been established to allow treated produce to be used for human consumption. TMRLs has been established for DIFENOCONAZOLE in CHARD [SILVERBEET], CHICORY LEAVES (GREEN AND RED CULTIVARS), ENDIVE AND SPINACH. MRLs can be found in the *Agricultural and Veterinary Chemicals Code Instrument No. 4 (MRL Standard)*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, residues must not exceed the limits/tolerances of the importing country.

Issued by the Australian Pesticides and Veterinary Medicines Authority.