



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF SCLEROTINIA STEM ROT IN LUPINS**

PERMIT NUMBER – PER91123

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 27 OCTOBER 2021 TO 31 OCTOBER 2024

Permit Holder:

PULSE AUSTRALIA LIMITED
Suite 501, Level 5
6 Help Street
CHATSWOOD NSW 2067

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

AVIATOR XPRO FOLIAR FUNGICIDE (APVMA No. 69361)

Containing: 150 g/L PROTHIOCONAZOLE and 75 g/L BIXAFEN as the only active constituents.

WARNING: Product may cause birth defects.

RESTRAINT:

DO NOT apply more than 2 applications per crop.

DO NOT apply after late flowering (BBCH69).

Adhere to label instructions on spray drift restraints and mandatory no-spray zones.

Directions for Use:

Crop	Disease	Rate
Lupins - Narrow leaf lupin, Albus lupin	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	550-800 mL/ha

Critical Use Comments:

- Apply initial foliar application during early flowering.
- Use higher rate when conditions are highly conducive for disease development.
- Apply using ground or aerial equipment in accordance with label instructions.
- Use a minimum spray volume of 70-100 L/ha for ground application, and 20 L/ha for aerial application.
- Apply a second application no sooner than 21 days later if disease persists, or if weather conditions are favourable for disease development.

Withholding Period:

Harvest: Not required when used as directed.

Grazing: Do not graze or cut for stock food for 35 days after application.

Livestock Destined For Export Markets

The grazing withholding periods apply to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval (ESI) is observed before stock are sold or slaughtered.

Export Slaughter Interval (ESI) – 9 Days

Livestock that have been grazing on treated crops should be placed on clean feed for 9 days prior to export slaughter.

Jurisdiction:

NSW only.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

Slight discolouration or necrosis on leaves ranging from specks to larger areas may occur on a small number of leaves per plant within two weeks of application. Any crop effects are likely to be generally negligible and not have any impact on grain yield. However, the sensitivity of the crop to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.

Export of treated produce

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with Aviator Xpro. If you are growing produce for export, please check with the permit holder for the latest information on MRLs and import tolerances before using Aviator Xpro.

Issued by the Australian Pesticides and Veterinary Medicines Authority