



PERMIT TO ALLOW MINOR USE OF AN AGVET CHEMICAL PRODUCT

PERMIT NUMBER -PER9187

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 30 SEPTEMBER 2007 TO 1 JANUARY 2010.

Permit Holder:

AUSVEG

C/- AgAware Consulting Pty Ltd

21 Rosella Avenue

STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Product to be used:

AMISTAR WG FUNGICIDE

AMISTAR 250 SC FUNGICIDE

Containing: 500 g/kg AZOXYSTROBIN

250 g/L AZOXYSTROBIN as the only active constituent.

Directions for Use:

Crop	Pest	Rate
PISTACHIO NUTS (<i>Pistacia vera</i>)	ALTERNARIA LATE BLIGHT (<i>Alternaria alternata</i>), <i>Botryosphaeria dothidea</i>	<u>For Amistar WG Fungicide</u> (500 g/kg Azoxystrobin): Apply 500 g/ha
	AND ANTHRACNOSE (<i>Colletotrichum acutatum</i>)	<u>For Amistar 250 SC Fungicide</u> (250 g/L Azoxystrobin): Apply 1 L/ha

Critical Use Comments:

Apply using orchard airblast/mister sprayer applying sufficient water to obtain uniform coverage. May be applied as a Dilute or Concentrate spray. DO NOT apply more than three times per season. Alternate with sprays of other chemical groups.

Dilute application: Water volumes typically range from 1800-20000 L/ha.

Concentrate application: Apply in 800-1000 L/ha.

Alternaria - Apply during nut development.

Anthracoze - Apply when conditions favour disease development.

Withholding Period:

DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

Jurisdiction:

ALL States

Additional Conditions:

THIS PERMIT provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

PERSONS who wish to prepare for use and/or use the products for the purposes specified in this permit must read, or have read to them, the permit particularly the information included in **DETAILS OF PERMIT** and **CONDITIONS OF PERMIT**.

Azoxystrobin is characterised by high toxicity to aquatic organisms. When applying Amistar, maintain a minimum buffer zone between treated pistachio trees and nearby waterbodies of 15 metres early in the season when there is not full leaf cover, and 10 metres late season when there is increased or full leaf cover.

TO AVOID CROP DAMAGE:

The sensitivity of some species and varieties of the crops to be treated under this permit has not been fully evaluated. It is advisable, therefore, to only treat a small number of plants to ascertain their reaction before treating the whole crop.

RESIDUES:

To allow produce from treated plants or animals to be supplied or otherwise made available for human or animal consumption the NRA has established the following temporary maximum residue limits (TMRLs): Azoxystrobin Tree nuts T0.02 mg/kg.

EXPORT OF PRODUCE:

Exported produce or animals must have appropriate residue tolerance limits established in the importing countries and any residues must not exceed the tolerance limits.

Issued by

Delegated Officer