



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF SCLEROTINIA ROT IN ADZUKI BEANS**

PERMIT NUMBER – PER82124

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 19 FEBRUARY 2016 TO 28 FEBRUARY 2025.

Permit Holder:

PULSE AUSTRALIA LIMITED
Suite 501, Level 5, 6 Help Street
CHATSWOOD NSW 2067

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

FILAN FUNGICIDE (APVMA No. 56362)

PLUS OTHER REGISTERED PRODUCTS

Containing: 500 g/kg BOSCALID as the only active constituent.

RESTRAINT:

Comply with all restraints on the product label.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- For maximum release height above the target canopy of 3 meters or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for aircraft')

Buffer zones for aircraft

Type of aircraft	Mandatory downwind buffer zones
	Natural aquatic areas
Fixed-wing	10 metres
Helicopter	15 metres

Directions for Use:

Crop	Pest / Disease	Rate
Adzuki Beans	Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i>)	0.775 – 1.0 kg product/ha

Critical Use Comments:

- Apply at 20-50% flowering to control white mould. Apply a second application 7-14 days later if disease persists, or weather conditions are favourable for disease development.
- Apply in a minimum spray volume of 100 L/ha for ground application or 40 L/ha for aerial application.
- DO NOT apply more than 2 applications per season.
- DO NOT apply less than 7 days after the initial treatment.

Re-entry period:

DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and the wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

Withholding Period:

Harvest: DO NOT harvest for 21 days after application.

Grazing: DO NOT graze or cut for stock food for 21 days after application.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that, **in addition** to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI) – 14 DAYS

LIVESTOCK THAT HAVE BEEN GRAZED ON, OR FED TREATED FORAGE, FODDER OR STUBBLE SHOULD BE PLACED ON CLEAN FEED FOR 14 DAYS PRIOR TO EXPORT SLAUGHTER.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

Export of treated produce

Temporary Maximum Residue Limits (TMRLs) have been established to allow treated produce to be used for human consumption. An MRL has been established for BOSCALID in ADZUKI BEAN (DRY). MRLs can be found in the *Agricultural and Veterinary Chemicals Code (MRL Standard) Instrument 2019*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, residues must not exceed the limits/tolerances of the importing country.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note: 14/08/2018 – Expiry date extended to 31/03/2022. Issued as version 2.

Note: 24/02/2022 – Permit updated to include spray drift restraints. Conditions of use updated to current standard.

Holder address updated. Permit expiry extended to 28/02/2025. Permit issued as Version 3.