



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF VARIOUS FUNGAL DISEASES IN NURSERY STOCK
(NON-FOOD)**

PERMIT NUMBER – PER88657

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 26 FEBRUARY 2020 TO 28 FEBRUARY 2023

Permit Holder:

NURSERY AND GARDEN INDUSTRY AUSTRALIA
C/o AgAware Consulting Pty Ltd
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

BANROT 400WP BROAD SPECTRUM FUNGICIDE FOR ORNAMENTALS (APVMA No. 52741)

Containing: 250 g/kg THIOPHANATE-METHYL and 150 g/kg ETRIDIAZOLE as the only active constituents.

BANROT 80G BROAD SPECTRUM FUNGICIDE FOR ORNAMENTALS (APVMA No. 53163)

Containing: 50 g/kg THIOPHANATE-METHYL and 30 g/kg ETRIDIAZOLE as the only active constituents.

Restrains:

DO NOT use on plants grown for human or animal consumption.

DO NOT use in the home garden.

Directions for Use:

BANROT 400WP BROAD SPECTRUM FUNGICIDE FOR ORNAMENTALS (APVMA No. 52741)

Crop	Disease	Rate	Critical Comments
Nursery stock (non-food): seedlings, tubes and plugs, potted colour, trees and shrubs, foliage plants, palms, grasses, and fruiting plants (non-bearing), cut flowers and ornamentals.	Damping off and root and stem rot diseases caused by the following fungi - <i>Pythium</i> , <i>Phytophthora</i> , <i>Rhizoctonia</i> , <i>Theilaviopsis</i> (<i>Chalara</i>)	<u>Pre-plant soil mix additive</u> 60 g / cubic metre	<ul style="list-style-type: none"> • Apply in accordance with approved product label directions. • Preventative treatment only and needs to be applied as close to infections events as possible. • Apply using a commercial soil/potting mixer. • DO NOT apply more than 4 applications per crop, with a minimum treatment interval of 4-8 weeks. • Follow the fungicide resistance warning on the product label. • Refer to product label for <i>Re-entry Period, Precautions, Safety Directions and First Aid Instructions.</i>
		<u>Post-plant soil drench</u> 400-800 g / 100 square metres in approximately 1000 L of water	<ul style="list-style-type: none"> • Apply in accordance with approved product label directions. • Apply using calibrated soil drench equipment. • DO NOT apply more than 4 applications per crop, with a minimum treatment interval of 4-8 weeks. • Use the lower rate when lower disease pressure is expected and the higher rate for higher disease pressure. • Follow the fungicide resistance warning on the product label. • Refer to product label for <i>Re-entry Period, Precautions, Safety Directions and First Aid Instructions.</i>

BANROT 80G BROAD SPECTRUM FUNGICIDE FOR ORNAMENTALS (APVMA No. 53163)

Crop	Disease	Rate	Critical Comments
Nursery stock (non-food): seedlings, tubes and plugs, potted colour, trees and shrubs, foliage plants, palms, grasses, and fruiting plants (non-bearing), cut flowers and ornamentals.	Damping off and root and stem rot diseases caused by the following fungi – <i>Pythium</i> , <i>Phytophthora</i> , <i>Rhizoctonia</i> , <i>Theilaviopsis</i> (<i>Chalara</i>)	<u>Pre-plant soil mix additive</u> 300 g / cubic metre	<ul style="list-style-type: none"> • Apply in accordance with approved product label directions. • Preventative treatment only and needs to be applied as close to infections events as possible. • Apply using a commercial soil/potting mixer. • DO NOT apply more than 4 applications per crop, with a minimum treatment interval of 4-8 weeks. • Refer to product label for <i>Re-entry Period, Precautions, Safety Directions and First Aid Instructions.</i>
		<u>Post-plant broadcast or pre-plant treatment</u> 2-4 kg / 100 square metres	<ul style="list-style-type: none"> • Apply in accordance with approved product label directions. • Apply using calibrated granule spreader. • DO NOT apply more than 4 applications per crop, with a minimum treatment interval of 4-8 weeks. • Use the lower rate when lower disease pressure is expected and the higher rate for higher disease pressure. • Refer to product label for <i>Re-entry Period, Precautions, Safety Directions and First Aid Instructions.</i>

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

The sensitivity of all nursery stock species and cultivars to Banrot when applied as a soil mix, broadcast or soil drench have not been fully evaluated under all growing conditions. It is advisable, therefore, to only treat a small area of crop to ascertain whether any phytotoxic reaction occurs before applying the spray treatment to the whole of a crop. Over-dosage may result in toxicity to sensitive plants. Do not continue treatment if the crop adversely reacts to the fungicide.