



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW SUPPLY AND MINOR USE OF AN UNREGISTERED AGVET
CHEMICAL**

PRODUCT FOR CONTROL OF SARCOPTIC MANGE IN WOMBATS

PERMIT NUMBER – PER89982

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a Supplier (as indicated) to possess the product for the purposes of supply and to supply the product to a person who can use the product under permit. This permit also allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder, the Supplier (if not one and the same) and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 14 DECEMBER 2020 TO 30 JUNE 2027

Permit Holder:

MANGE MANAGEMENT INCORPORATED
78 Caledonia Street
ST ANDREWS VIC 3761

Persons authorised for supply:

- The Permit Holder; and
- Wombat Mange Treatment Groups as authorised in writing by the Permit Holder.

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

CYDECTIN POUR-ON FOR CATTLE AND RED DEER (APVMA No. 45970)
PLUS OTHER REGISTERED PRODUCTS
Containing: 5 g/L MOXIDECTIN as the only active constituent and 150 g/L
HYDROCARBON LIQUID as the only labelled solvent.

OR

MOXIDECTIN FOR WOMBATS WITH MANGE

AN UNREGISTERED PRODUCT prepared by decanting:
CYDECTIN POUR-ON FOR CATTLE AND RED DEER (APVMA No. 45970)
OR OTHER REGISTERED PRODUCTS
Containing: 5 g/L MOXIDECTIN as the only active constituent and 150 g/L
HYDROCARBON LIQUID as the only labelled solvent.

RESTRAINT:

DO NOT use in conjunction with any other treatments.
DO NOT decant any product into drink bottles.
DO NOT treat any wombats unless trained in the identification, classification the stages of mange infection and observation of toxicity signs in wombats.

Directions for Use:

Host	Pest	Rate
Wombats	Sarcoptic mange (<i>Sarcoptes scabiei</i>)	0.8ml per kg body weight (maximum dose of 20mL on an adult wombat)

Critical Use Comments:

- Apply once weekly for 15 weeks, delivered via a burrow flap or direct application with a pole & scoop to the wombat's back.
- Apply along the backline of the wombat using the Pole and Scoop method. A pole is, for instance, a broom or mop handle with a scoop firmly attached to the end which is able to hold the appropriate dose.
- Burrow flap application may be used when wombats cannot be approached to successfully apply the treatment using pole and scoop. E.g. wombats which are recovering and run away when approached.
- DO NOT split doses when applying by burrow flap. The full dose for each single treatment must be applied at one time, or there is a risk of underdosing, and contributing to resistance development.
- The product must only be used by individuals who have been trained in classifying the stages of mange infection and observation of toxicity signs.
- Continued treatment in wombats which showed toxicity signs after the first treatment should be stopped or the dose reduced in consultation with a registered veterinarian.
- Wombats with visible lacerations along the backline should not be treated with moxidectin.
- DO NOT overdose animals. Due to the lipophilic nature of moxidectin and the lower proportion of body fat carried by wombats, an acute (immediate) poisoning effect is more likely to be seen in a wombat than chronic (long term) poisoning by repeated dosing over a long period.

- A sign as detailed in **Attachment 2** must be placed in areas where wombats are undergoing treatment. The phone number of the person applying the treatment to the wombat must be written in the relevant space on the sign.

First Aid Directions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

Safety Directions:

Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Do not inhale. May irritate the nose and throat.

Jurisdiction:

All States and Territories.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the *Agricultural and Veterinary Chemicals Code Act 1994*, in accordance with the obligation imposed by that section.

Supply

The supplier must supply the product in a container that complies with the requirements of section 18 of the Agricultural and Veterinary Chemicals Code Regulations.

The registered product may be decanted into smaller containers (up to 1L poison bottles with a child proof lid), this product is then considered unregistered, and is referred to as 'Moxidectin for Wombats with Mange'. Decanted product must only be supplied from containers with more than 12 months left of the expiry date.

Attached to this container must be a label which is identical in content and format to the label at **Attachment 1**.

Moxidectin for Wombats with Mange must only be supplied as part of a wombat mange treatment kit, which must contain at minimum:

- a) A copy of the label for the registered product which has been supplied in the smaller container,
- b) a printout of the sign consistent in content with the text in **Attachment 2** and instructions on its placement,
- c) a copy of this permit. Any additional directions for use must be consistent with this permit.

Where treaters are using Moxidectin for Wombats with Mange, the decanted amount must be used immediately within 1 treatment cycle (15 weeks) in 1-2 wombats.

DO NOT supply in drink bottles.

Target Animal Safety and Efficacy

The efficacy and toxicity associated with this dose rate has not been fully evaluated. Users must observe wombats for any toxicity signs as detailed in the critical use comments.

Users must also monitor animals for the efficacy of the Product, and information on the efficacy or any lack of efficacy provided to the permit holder.

Reporting

The Permit Holder must record any adverse experience associated with the use of the product that they are made aware of and report them to the APVMA as soon as possible, ph. 1800 700 583 or aerp@apvma.gov.au.

In addition to reporting to the APVMA, any wombats that die after treatment should be reported to a veterinarian and a detailed post mortem should be carried out.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note: 15/12/2023 – Permit updated to update conditions to current template wording, to add AERP phone number, and to update container requirements, use of decanted product and restraints. Permit expiry extended to 30/06/2024. Permit issued as Version 2.

12/09/2024 – Permit updated to clarify directions and bring it in line with PER89040 where appropriate. Permit expiry extended to 30/06/2027. Permit issued as Version 3.

Attachment 1: Label

**CAUTION
FOR ANIMAL TREATMENT ONLY
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

MOXIDECTIN FOR WOMBATS WITH MANGE

**5g/L Moxidectin
150 g/L Hydrocarbon Liquid solvent**

Dose rate: 0.8ml / 1kg wombat body weight

User MUST Read APVMA Permit 89982 before use

Poisons Information Centre. Phone: 13 11 26
www.mangemanagement.org.au

Date of decanting: _____

WOMBAT MANGE TREATMENT

A wombat with mange lives in this area.

The wombat is being treated using a treatment that is applied to its skin via a burrow flap device or directly to its back via a pole and scoop (APVMA Permit 89982).

If you see this wombat, please stay away from it.

WARNING: Do not touch or interfere with the burrow flap device

The product that has been used contains: 5 g/L Moxidectin as the active, and 150 g/L Hydrocarbon Liquid solvent.

First Aid Directions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

Safety Directions:

Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Do not inhale. May irritate the nose and throat.

For more information contact the permit holder:

www.mangemanagement.org.au

Phone: 0431 600 125

E: info@mangemanagement.org.au

Phone number for the person applying the treatment: _____