



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF ANTHRACNOSE IN NON-FOOD NURSERY STOCK**

PERMIT NUMBER – PER90451

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 25 OCTOBER 2021 TO 31 OCTOBER 2024

Permit Holder:

GREENLIFE INDUSTRY AUSTRALIA LIMITED
C/o – AgAware Consulting Pty Ltd
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

OCTAVE WP FUNGICIDE (APVMA No. 30485)

PLUS OTHER REGISTERED PRODUCTS

Containing: 462 g/kg PROCHLORAZ AS THE MANGANESE CHLORIDE COMPLEX as the only active constituent.

SPORTAK FUNGICIDE (APVMA No. 30484)

PLUS OTHER REGISTERED PRODUCTS

Containing: 450 g/L PROCHLORAZ as the only active constituent.

RESTRAINT:

DO NOT use on vegetable and herb seedlings.

DO NOT feed treated crops to livestock, or allow livestock to graze treated areas.

DO NOT use on crops destined for export.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply by ground boomspray unless the following mandatory buffer zones are observed:

Droplet size	Bystander buffer zone (metres)
Fine	10
Medium	0

Directions for Use:

Crop	Disease	Rate
Nursery Stock (Non-Food)	Anthracnose <i>(Colletotrichum spp)</i>	100 g product / 100 L Or
Seedlings & plugs (not including vegetable and herb seedlings) Potted colour Trees & shrubs Foliage plants Palms Grasses Fruit plants (non-bearing)		100 mL product / 100 L

Critical Use Comments:

- Prochloraz based products should be applied as a protectant treatment prior to the onset of disease development.
- Apply initial application at the first signs of disease infection. If conditions remain favourable, repeat application at 7 – 14 day interval.
- Apply maximum of three (3) foliar spray applications per crop.
- Apply using calibrated air-blast sprayer, air-shear sprayer, back-pack mister, hand-lance sprayer, boom sprayer or similar equipment.
- Apply in sufficient water volume to ensure thorough coverage of all plant surfaces. Avoid excessive runoff of diluent onto the ground.
- DO NOT allow spray mix to stand overnight.
- DO NOT apply foliar treatment if rainfall is expected or irrigation is scheduled before the spray is dry.

Jurisdiction:

All States and Territories.

Safety Directions:

Product is poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin.

When using together with other products, consult their label safety directions.

When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). In addition, when using the prepared spray by hand held equipment wear elbow-length chemical resistant gloves.

If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water.

Wash hands after use. After each day's use, wash gloves, face shield and contaminated clothing.

Re-Entry:*General Public –*

DO NOT allow entry into treated areas until the spray has dried.

Occupational Users –

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

For glasshouses and other confined areas, do not re-enter until spray deposits have dried and area has been thoroughly ventilated.

If performing hand-set irrigation activities, DO NOT enter treated areas for three days unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

The sensitivity of the crop to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.

Issued by the Australian Pesticides and Veterinary Medicines Authority