



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW SUPPLY AND EMERGENCY USE OF AN UNREGISTERED AGVET
CHEMICAL PRODUCT FOR CONTROL OF MICE IN CERTAIN CROPS AND PASTURE**

PERMIT NUMBER – PER90799

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a Supplier (as indicated) to possess the product for the purposes of supply and to supply the product to a person who can use the product under permit. This permit also allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder, the Supplier (if not one and the same) and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 7 MAY 2021 TO 31 OCTOBER 2022

Permit Holder:

GRAIN PRODUCERS AUSTRALIA LIMITED
13 Wallace Street
BRAIDWOOD NSW 2622

Suppliers:

ANIMAL CONTROL TECHNOLOGIES (AUSTRALIA) PTY LTD
Street 46-50 Freight Drive
SOMERTON VIC 3062

IMTRADE AUSTRALIA PTY LTD
Suite 22, 11 Preston Street
COMO WA 6152

PCT HOLDINGS PTY LTD
Unit 5, 47 Murdoch Circuit
ACACIA RIDGE QLD 4110

WILHELM RURAL PTY LTD
116 Sixth Avenue
JOSLIN SA 5070

4 FARMERS AUSTRALIA PTY LTD
35a Mcdowell Street
WELSHPOOL WA 6106

TRIOX PTY LTD
6,707 Mt Alexander Rd
MOONEE PONDS VIC 3039

Persons who can use the product under this permit:

Persons authorised or licenced under the relevant state or territory poisons control for Schedule 7 Poisons.

CONDITIONS OF USE

Products to be used:

MOUSEOFF 50 ZINC PHOSPHIDE BAIT

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

IMTRADE DEADMOUSE DOUBLE STRENGTH ZINC PHOSPHIDE BAIT

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

SUREFIRE 50 ZINC PHOSPHIDE MOUSE BAIT

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

SUREFIRE 50 ZINC PHOSPHIDE MOUSE BAIT PELLETS

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

LAST SUPPER SUPREME 50 ZINC PHOSPHIDE MOUSE BAIT

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

LAST SUPPER SUPREME 50 ZINC PHOSPHIDE MOUSE BAIT PELLETS

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

4 FARMERS ZINC PHOSPHIDE 50G/KG UNSTERILIZED MOUSE BAIT

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent.

4 FARMERS ZINC PHOSPHIDE PELLET

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent

PESTMASTER ZINC PHOSPHIDE MOUSE KILLER PELLETS

AN UNREGISTERED PRODUCT

Containing: 50 g/kg ZINC PHOSPHIDE as the only active constituent

RESTRAINTS:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crops must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

DO NOT apply if heavy rain is imminent.

Directions for Use:

Crop	Pest	Rate
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations only a post-emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).	Mice (<i>Mus musculus</i>)	1 kg / ha to achieve an even coverage of 2 - 3 grains or pellets / m ²

Critical Use Comments:

- Apply when mice reach threshold when severe damage is being caused to broadacre crops.
- Apply using ground or aerial equipment according to the label directions.
- Apply bait only when pre-baiting monitoring by bait cards indicates high population presence only.
- Apply only at night or early evening to minimise non-target primary consumption and maximise bait consumption by mice.
- Ensure that there is adequate ground cover (e.g. crop stubble, crop, pasture etc.) to minimise primary bait consumption
- Apply application evenly across the treatment area. DO NOT leave piles of bait in treatment area as this will increase opportunistic feeding by non-target birds, kangaroo, wallabies, livestock etc.
- Users need to consider and familiarise themselves with baiting strategies to ensure greater success for treatment site (e.g. co-ordinated bait timing with neighbours, bait adjacent areas etc.) Refer to GRDC ‘Tips and Tactics Better Mouse Management’ or similar relevant document from state department of Agriculture.
https://grdc.com.au/__data/assets/pdf_file/0020/243803/GRDC-Tips-and-Tactics-Better-Mouse-Management-National-2017-low-res.pdf?utm_source=website&utm_medium=download_link&utm_campaign=pdf_download&utm_term=National&utm_content=Tips%20and%20Tactics:%20Better%20Mouse%20Management
- Eliminate as far as practicable all alternative food sources to increase bait desirability and consumption at treatment site.
- Repeat treatments will likely be required when high populations are observed. Continue to monitor site with bait cards to determine population size, application rate and if re-treatment is required.
- For all other directions, refer to the appropriate labels at **Attachments 1 through 9**.

Withholding Period:

Harvest: Do not harvest for 14 days after application.

Grazing: Do not graze or cut for stock food for 14 days after application.

Jurisdiction:

All States and Territories.

Safety Directions for Loading Bait for Aerial Application:

When opening container and loading the bait, wear elbow-length PVC gloves and full face piece respirator with combined dust and gas cartridge (or supplied air respirator).

Protection of Wildlife, Fish, Crustaceans and Environment:

Toxic to native wildlife and birds. Fields must be closely monitored for non-target species during a pre-baiting period. DO NOT apply if non-target animals are found feeding during the pre-baiting period. DO NOT use on sites where vulnerable, threatened, endangered or critically endangered mammals or birds are known to occur.

Any incidents where it is suspected that non-target animals may have been poisoned should be reported to State or Territory authorities.

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Disposal:

To the extent possible, animal carcasses should be recovered during and for 14 days after a baiting campaign and be destroyed by burning or burial according to the requirements of the State or Territory in which use has occurred.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the appropriate product label at **Attachments 1 through 9**.

Supply

The supplier must supply the product in a container that complies with the requirements of section 18 of the Agricultural and Veterinary Chemicals Code Regulations. Attached to this container must be a label which is identical in content and format to the appropriate label at **Attachments 1 through 9**.

Record keeping

Maintain records of all treatments performed under this permit. Details must include the date and locations where treatments occurred, the total amount of product used, and the names and addresses of the persons undertaking the use. These details must be maintained for a minimum period of two years from the date of expiry of this permit, and must be made available to the APVMA upon request.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note: 6/10/2021. Permit updated to add two products, a supplier and extend the permit to 31/10/2022. Permit issued as Version 3

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**MOUSEOFF 50
ZINC PHOSPHIDE BAIT**

**ACTIVE CONSTITUENTS:
50 g/kg ZINC PHOSPHIDE**

**For the control of heavy infestations of mice around commercial and industrial premises and in
agricultural situations as per the directions of use**

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROLS FOR SCHEDULE 7 POISONS.**

Net Contents: 5 – 500 kg

**THIS PRODUCT IS NOT REGISTERED.
Product is to be used in accordance with relevant APVMA permit – PER90799**

**Animal Control Technologies (Australia) Pty Ltd
46-50 Freight Drive, Somerton VIC 3062
www.animalcontrol.com.au
Telephone Contact 03 9308 9688**

DIRECTIONS FOR USE

Restraints

DO NOT apply bait to bare ground (Including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

SITUATION	STATE	PEST	RATE	CRITICAL COMMENTS
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations-only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).	All	Mice	1 kg/ha to achieve an even coverage of 2-3 grains/m ²	Bait only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m ² . It is imperative that all attempts are made to apply bait as evenly as possible to maximize product efficacy and minimize risks to non- target animals. MOUSEOFF 50 Zinc Phosphide Bait will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment after 14 days may therefore be required in certain circumstances. It is recommended pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Dispose of use containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames- forms toxic gas. When opening the container and using the product, wear elbow-length PVC gloves. and a full facepiece respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use, wash gloves and respirator and if rubber, wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (phone: 13 11 26). Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air- viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet.

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY:

This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the Directions for Use given on the label. No responsibility is accepted in respect of this product save those non-excludable conditions implied by any Federal or State legislation or law of a Territory.

BN:

DOM:

DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE

DEADMOUSE

DOUBLE STRENGTH
ZINC PHOSPHIDE BAIT

ACTIVE CONSTITUENT: 50g/kg ZINC PHOSPHIDE



For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR
LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR
SCHEDULE 7 POISONS.

IMPORTANT: READ THIS LABEL THOROUGHLY BEFORE OPENING
OR USING THIS PRODUCT

**FOR USE ONLY UNDER EMERGENCY
PERMIT NO. PER90799**



17 Ocean St, Kwinana Beach WA 6167
(08) 9419 0333
www.imtrade.com.au

**DIRECTIONS FOR USE
RESTRAINTS**

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

Situation	Pests	Rate	Critical Comments
<p>Grain crops, Legume crops, Canola, Safflower, Nut crops and Pasture.</p> <p>(Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).</p>	Mice	1kg/ha to achieve an even coverage of 2-3 grains/m ²	<p>Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m².</p> <p>It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals.</p> <p>Imtrade DeadMouse Zinc Phosphide Bait will provide a high level of control in most situations. However, in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances.</p> <p>It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control.</p> <p>Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

**THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN.
DO NOT USE THIS PRODUCT IN THE HOME GARDEN.**

WITHHOLDING PERIOD

DO NOT HARVEST ANY CROPS OR ALLOW LIVESTOCK TO GRAZE BAITED AREAS FOR 14 DAYS AFTER APPLICATION.

PROTECTION OF WILDLIFE, LIVESTOCK AND THE ENVIRONMENT

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Dispose of used containers by crushing and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging one meter below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. **DO NOT** inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits, wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). **DO NOT** give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

Additional GHS Hazard & Precautionary Statements

●In contact with water releases flammable gases which may ignite spontaneously; ●Very toxic to aquatic life with long lasting effects; ●DO NOT eat, drink or smoke when using this product; ●IF SWALLOWED: Rinse mouth. Do NOT induce vomiting; ●If exposed or if you feel unwell: Call a POISON CENTER or doctor/physician; ●Collect spillage; ●Not combustible. Use extinguishing media suited to burning materials (NOT WATER).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

In a Transport Emergency
Dial 000
Police or Fire Brigade

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**SureFire® 50
ZINC PHOSPHIDE MOUSE BAIT**

ACTIVE CONSTITUENT: 50 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROLS FOR SCHEDULE 7 POISONS.**

Formula B Unsterilised grain only

**RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM
UNSTERILISED GRAIN**

This product may have been prepared using unsterilized wheat grain that has not been subjected to sterilization by irradiation as indicated below. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilized bait products are used.

PRODUCT MADE USING UNSTERILISED GRAIN.
UNSTERILISED GRAIN SOURCED FROM: _____

NET CONTENTS: 5 kg - 500 kg

PCT Holdings Pty Ltd

® Surefire is a registered trademark of PCT International Pty Ltd
(PCT Holdings Pty Ltd ABN 11 099 023 962)

5/74 Murdoch Circuit

ACACIA RIDGE QLD 4110

<http://pct.au.com>

CUSTOMER SERVICE FREECALL 1800 630 877

EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**FOR USE UNDER PERMIT 90799
THIS PRODUCT IS NOT REGISTERED**

DIRECTIONS FOR USE

Restraints:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

Situation	Pest	Rate kg/ha	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover).	Mouse	1 kg/ha to achieve an even coverage of 2-3 grains/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m ² . It is imperative that all attempts are made to bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Surefire 50 Zinc Phosphide Mouse Bait will provide a high level of control in most situations. However in extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS

THIS PRODUCT IS TOO DANGEROUS TO BE USED IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD: DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

GENERAL INSTRUCTIONS

RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM UNSTERILISED GRAIN

This product may be prepared using unsterilised wheat grain that has not been subjected to sterilisation by irradiation. The main panel of the label of this product indicates whether this product has been prepared using unsterilised grain. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilised bait products are used.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container. Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of empty containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which is available from PCT Holdings Pty Ltd on request. Call Customer Service Toll Free on 1800 630 877 or visit our web site at <http://pct.au.com>

NOTICE

PCT Holdings Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by PCT Holdings Pty Ltd, or under abnormal conditions.

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**SureFire® 50
ZINC PHOSPHIDE MOUSE BAIT PELLETS**

ACTIVE CONSTITUENT: 50 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROLS FOR SCHEDULE 7 POISONS.**

NET CONTENTS: 5 kg - 500 kg

PCT Holdings Pty Ltd

® Surefire is a registered trademark of PCT International Pty Ltd
(PCT Holdings Pty Ltd ABN 11 099 023 962)
5/74 Murdoch Circuit
ACACIA RIDGE QLD 4110
<http://pct.au.com>

*CUSTOMER SERVICE FREECALL 1800 630 877
EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877*

**FOR USE UNDER PERMIT 90799
THIS PRODUCT IS NOT REGISTERED**

DIRECTIONS FOR USE

Restraints:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

Situation	Pest	Rate kg/ha	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover).	Mouse	1 kg/ha to achieve an even coverage of 2-3 pellets/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 pellets/m ² . It is imperative that all attempts are made to bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Surefire 50 Zinc Phosphide Mouse Bait Pellets will provide a high level of control in most situations. However in extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS. THIS PRODUCT IS TOO DANGEROUS TO BE USED IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD: DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container. Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of empty containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which is available from PCT Holdings Pty Ltd on request. Call Customer Service Toll Free on 1800 630 877 or visit our web site at <http://pct.au.com>

NOTICE

PCT Holdings Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by PCT Holdings Pty Ltd, or under abnormal conditions.

Batch No:

Date of Manufacture:

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Last Supper
Supreme 50
Zinc Phosphide Mouse Bait**

ACTIVE CONSTITUENT: 50 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROL FOR SCHEDULE 7 POISONS.**

NET CONTENTS: 5 kg - 500 kg.

Formula B Unsterilised grain only

**RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM
UNSTERILISED GRAIN**

This product may have been prepared using unsterilized wheat grain that has not been subjected to sterilization by irradiation as indicated below. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilized bait products are used.

PRODUCT MADE USING UNSTERILISED GRAIN.

UNSTERILISED GRAIN SOURCED FROM: _____

Produced by:

Wilhelm Rural Pty Ltd

ABN 22 055 833 111

116 Sixth Ave, Joslin South Australia 5070

Ph: 0419 821 422

Email: bwilhelm@coburnwilhelm.com

**FOR USE UNDER PERMIT 90799
THIS PRODUCT IS NOT REGISTERED**

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

SITUATION	PEST	RATE	CRITICAL COMMENTS
<p>Grain crops, legume crops, canola, safflower, nut crops and pasture.</p> <p>(Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover)</p>	Mice	1 kg/ha to achieve an even coverage of 2 – 3 grains/m ²	<p>Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2 - 3 grains/m². It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Last Supper Supreme 50 Zinc Phosphide Mouse Bait will provide a high level of control in most situations. However in extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances.</p> <p>It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control.</p> <p>Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT HARVEST ANY CROPS OR ALLOW LIVESTOCK TO GRAZE BAITED AREAS FOR 14 DAYS AFTER APPLICATION.

RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM UNSTERILISED GRAIN

This product may be prepared using unsterilised wheat grain that has not been subjected to sterilisation by irradiation. The main panel of the label of this product indicates whether this product has been prepared using unsterilised grain. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilised bait products are used.

GENERAL INSTRUCTIONS

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of empty containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames — forms toxic gas. When opening the container and using the product wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

SAFETY DATA SHEET

Additional information is listed in the SDS which can be obtained from the supplier.

CONDITIONS OF SALE

The use of Last Supper Zinc Phosphide Supreme 50 Mouse Bait being beyond the control of the manufacturer, no warranty expressed or implied is given by Wilhelm Rural Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wilhelm Rural Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

BN: DOM:

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Last Supper
Supreme 50
Zinc Phosphide Mouse Bait Pellets**

ACTIVE CONSTITUENT: 50 g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROL FOR SCHEDULE 7 POISONS.**

Net Contents: 5 kg - 500 kg

Wilhelm Rural Pty Ltd
ABN 22 055 833 111
116 Sixth Avenue Joslin SA 5070

Phone: 0419 821 422
Email: bwilhelm@coburnwilhelm.com

**FOR USE UNDER PERMIT 90799
THIS PRODUCT IS NOT REGISTERED**

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crops must be employed to reduce risk to birds and other non-target animals.

DO NOT apply unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed has been sown into crop stubble or other similar ground cover).	Mice	1 kg/ha to achieve an even coverage of 2 – 3 pellets/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2 – 3 pellets/m ² . It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. This product will provide a high level of control in most situations. However, in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying the bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

**THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN.
DO NOT USE THIS PRODUCT IN THE HOME GARDEN.**

WITHHOLDING PERIOD

DO NOT HARVEST ANY CROPS OR ALLOW LIVESTOCK TO GRAZE BAITED AREAS FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of empty containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using the product wear elbow length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

SAFETY DATA SHEET

Additional information is listed in the SDS which can be contained by the supplier.

CONDITIONS OF SALE

The use of Last Supper Supreme 50 Zinc Phosphide Mouse Bait Pellets being beyond the control of the manufacturer, no warranty expressed or implied is given by Wilhelm Rural Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wilhelm Rural Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

BN: DOM:

4 FARMERS ZINC PHOSPHIDE 50g/kg UNSTERILIZED MOUSE BAIT

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ACTIVE CONSTITUENT: 50g/kg ZINC PHOSPHIDE

THIS PRODUCT IS NOT REGISTERED.

Product is to be used in accordance with relevant APVMA permit – PER90799

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

RISKS OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM UNSTERILISED GRAIN

This product 4Farmers Zinc Phosphide Unsterilised Mouse Bait is prepared using unsterilized wheat grain that has not been subjected to sterilisation by irradiation. Thus the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that the fullysterilised versions of Zinc Phosphide Mouse Baits are used.

UNSTERILISED GRAIN SOURCED FROM:

UNSTERILISED GRAIN

This product 4Farmers Zinc Phosphide Unsterilised Mouse Bait is prepared using unsterilized wheat grain that has not been subjected to sterilisation by irradiation. Thus the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that the fully sterilised versions of Zinc Phosphide Mouse Baits are used

Net Contents: 1-125kg
150kg to 500 kg

Restraints:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover)

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance

Directions for Use

Situation	Pest	Rate	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops, and pasture. (Note: for all situations – only as a post emergent treatment, stubbles of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover.	Mice	1 kg/ha to achieve an even coverage of 2 – 3 grains/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m ² . It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Farmers Zinc Phosphide Unsterilized Bait will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

Storage and Disposal:

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

Safety Directions:

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames - forms toxic gas. When opening the container and using baits wear elbow-length PVC gloves and a full-face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 1126). Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

4 FARMERS ZINC PHOSPHIDE 50g/kg UNSTERILIZED MOUSE PELLET

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ACTIVE CONSTITUENT: 50g/kg ZINC PHOSPHIDE

THIS PRODUCT IS NOT REGISTERED.

Product is to be used in accordance with relevant APVMA permit – PER90799

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROLS FOR SCHEDULE 7 POISONS.

Net Contents: 1-125kg
150kg to 500 kg

Restraints:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover)

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals

DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance

Directions for Use

Situation	Pest	Rate	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops, and pasture. (Note: for all situations – only as a post emergent treatment, stubbles of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover.	Mice	1 kg/ha to achieve an even coverage of 2 – 3 pellets/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 pellets/m ² . It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Farmers Zinc Phosphide 50g/kg Mouse Pellet will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Retreatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

Storage and Disposal:

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition.

Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

Safety Directions:

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames - forms toxic gas. When opening the container and using baits wear elbow-length PVC gloves and a full-face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use wash gloves and respirator and if rubber wash with detergent and warm water.

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 1126). Do not give direct mouth-to-mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one-way mask. Resuscitate in a well-ventilated area.

**DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**PESTMASTER
ZnP 50
MOUSE KILLER PELLETS**

ACTIVE CONSTITUENT: 50g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

**THIS PRODUCT IS NOT REGISTERED.
Product is to be used in accordance with relevant APVMA permit – PER90799**

**THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS
AUTHORIZED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY
POISONS CONTROLS FOR SCHEDULE 7 POISONS.**

Net Contents: 15kg - 500kg

RESTRAINTS:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

DIRECTIONS FOR USE

Situation	Pest	Rate	Critical Comments
<p>Grain crops, legume crops, canola, safflower, nut crops and pasture.</p> <p>(Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seed have been sown into crop stubble or other similar ground cover).</p>	Mice	1kg/ha to achieve an even coverage of 2-3 pellet/m ²	<p>Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 pellet/m². It is imperative that all attempts are made to apply bait as evenly as possible to maximize product efficacy and minimise risks to non-target animals.</p> <p>PESTMASTER ZnP 50 Mouse Killer Pellets will provide a high level of control in most situations. However in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances.</p> <p>It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying the bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control.</p> <p>Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.</p>

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISES UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD: DO NOT harvest any crops or allow livestock to graze baited areas for 14 days after application.

PROTECTION OF WILDLIFE, THE ENVIRONMENT AND LIVESTOCK

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

STORAGE AND DISPOSAL The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using baits wear elbow length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Do not give direct mouth to mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy viva or one way mask. Resuscitate in a well ventilated area. **SAFETY DATA SHEET** Additional information is listed in the SDS which can be obtained from the supplier.

NOTICE: Triox Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Triox Pty Ltd or under abnormal conditions.

Batch Number: DOM:

Triox Pty Ltd
6/707 Mount Alexander Road, Moonee Ponds VIC 3039
Information Line: 1300 729 354

EMERGENCY CONTACT NUMBER 1300 729 354