

Australian Pesticides and Veterinary Medicines Authority

PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL

PRODUCT FOR CONTROL OF PHOMOPSIS/DIAPORTHE DISEASE IN SOYA BEAN

SEEDS

PERMIT NUMBER – PER92352

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 26 OCTOBER 2022 TO 31 OCTOBER 2025

Permit Holder: AUSTRALIAN OILSEEDS FEDERATION INC 30 Coronga Crescent KILLARA NSW 2071

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used: MAXIM 100 FS FUNGICIDE SEED TREATMENT (APVMA No. 51407) PLUS OTHER REGISTERED PRODUCTS Containing: 100 g/L FLUDIOXONIL as the only active constituent.

Directions for Use:

Сгор	Disease	Rate
Soya Bean seeds for planting	Phomopsis/diaporthe disease (Diaporthe spp.)	50 mL / 100 kg seed

Critical Use Comments:

- Apply as a slurry before sowing at a rate of 5 10 mL of slurry per kg of seed.
- Apply to seed using commercial seed treatment equipment.
- DO NOT store treated seed for more that 6 months before sowing.
- DO NOT apply more than 1 application per seed.
- DO NOT use treated seed for animal or human consumption.
- DO NOT allow treated seed to contaminate other seed intended for animal or human consumption.
- When treated seed is stored, it should be kept apart from other seed and the bags or other containers should be clearly marked to indicate the contents have been treated with this product. Bags and containers which have held treated seed are not to be used for any other purpose.
- DO NOT feed treated seed, or otherwise expose, to wild or domestic birds. Any spillage of treated seed which occurs either during the seed treatment process or in the field operations must be cleaned up immediately, preferably by recovery and re-use.
- It is recommended not to use seed lots with more than 20 percent affected seed because severely infected or mouldy seed will fail to germinate even after being treated.

Withholding Period:

Harvest: Not required when used as directed. Grazing: Do not graze or cut for stock food for 7 weeks after application.

Protection of Livestock:

DO NOT feed treated seed to animals.

DO NOT allow seed treated with this product to contaminate feed intended for animal consumption.

DO NOT allow any feed material that has come in contact with this product to be fed to animals.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the *Agricultural and Veterinary Chemicals Code Act 1994*, in accordance with the obligation imposed by that section.

To Avoid Crop Damage

The sensitivity of the crop to be treated under this permit has not been fully evaluated. It is advisable to treat a small number of seeds and test them for successful germination before treating the whole crop.

Issued by the Australian Pesticides and Veterinary Medicines Authority