



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW EMERGENCY USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF GREY MILDEW IN COTTON**

PERMIT NUMBER – PER92749

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 5 SEPTEMBER 2022 TO 30 SEPTEMBER 2025

Permit Holder:

COTTON AUSTRALIA LTD
Suite 4.01, 247 Coward St
MASCOT NSW 2020

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

MAXENTIS EC FUNGICIDE (APVMA No. 89549)

Containing: 133 g/L AZOXYSTROBIN and 100 g/L PROTHIOCONAZOLE as the only active constituents.

RESTRAINT:

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff from treated fields for at least 3 days after application.

For PROFESSIONAL use only.

Directions for Use:

Crop	Disease	Rate
Cotton	Grey mildew (<i>Ramulariopsis</i> sp.)	300-900 mL/ha

Critical Use Comments:

- Apply at first sign of disease development.
- Apply by aerial or ground boomspray.
- Use a spray volume of at least 20 L/ha for aerial application or 70 L/ha for ground application.
- Apply higher rates for better disease control, longer residual activity and higher yield potential.
- Use Hasten wetter at 500 mL/100 L.
- DO NOT apply more than 2 applications per crop.
- DO NOT apply less than 14-21 days after the initial treatment.

Withholding Period:

Harvest: DO NOT harvest for 5 weeks after application.

Grazing: DO NOT allow livestock to graze treated cotton crop, stubble or gin trash.

Spray drift restraints:

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.

Follow all Spray Drift on the product label.

First Aid Warnings:

ADDITIONAL USER SAFETY INFORMATION WARNING: May cause birth defects.

Jurisdiction:

NT, Qld and WA only - locations above the 26 parallel line.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

The product label notes that treatment may result in minor phytotoxicity symptoms in canola crops i.e., bleaching or spotting of leaves. These effects are generally negligible and not expected to negatively impact grain yield. However, the sensitivity of the crop (cotton) to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.

Export of treated produce

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with azoxystrobin and prothioconazole in *Maxentis EC Fungicide*. If you are growing produce for export, please check with the permit holder for the latest information on MRLs and import tolerances before using azoxystrobin and prothioconazole.

Issued by the Australian Pesticides and Veterinary Medicines Authority