



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF VARIOUS INSECTS IN NURSERY STOCK**

PERMIT NUMBER – PER92919

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 6 OCTOBER 2023 TO 31 OCTOBER 2026

Permit Holder:

GREENLIFE INDUSTRY AUSTRALIA LIMITED
c/o AgAware Consulting Pty Ltd.
21 Rosella Avenue
STRATHFIELDSAYE VIC 3551

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

MINECTO FORTE INSECTICIDE (APVMA No. 87610)

Containing: 80 g/L CYANTRANILIPROLE + 400 g/L DIAFENTHIURON as the only active constituent.

RESTRAINT:

DO NOT apply by aircraft.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT apply consecutive applications.

DO NOT apply when honeybees are actively foraging.

SPRAY DRIFT RESTRAINTS

Read and follow all spray drift restraints on the product label.

Mandatory downwind buffer zones - Natural aquatic areas 5meters

Directions for Use:

Crop	Pest	Rate
Nursery stock (non-food) - seedlings & plugs, potted colour, trees & shrubs, foliage plants, palms, grasses, fruit plants (non-bearing), cut flowers and ornamentals (field and protected)	Silverleaf whitefly (<i>Bemisia tabaci</i>)	500 mL/ha or 50 mL/100 L
	Melon aphid (<i>Aphis gossypii</i>) Green peach aphid (<i>Myzus persicae</i>)	500-625 mL/ha or 50-62.5 mL/100 L
	<i>Helicoverpa</i> spp. Cucumber moth (<i>Diaphania indica</i>) Potato moth (<i>Phthorimaea operculella</i>) Cluster caterpillar (<i>Spodoptera litura</i>)	500 mL/ha or 50 mL/100 L
	Two-spotted mite (<i>Tetranychus urticae</i>)	750 mL/ha or 75 mL/100 L
	Tomato thrips (<i>Frankliniella schultzei</i>) Western flower thrips (<i>Frankliniella occidentalis</i>) Plague thrips (<i>Thrips imaginis</i>) (suppression only)	750 mL/ha or 75 mL/100 L

Critical Use Comments:

- Apply to newly developing infestations.
- Apply foliar spray by air-blast sprayer, boom sprayer or equivalent.
- Use sufficient spray volume to ensure coverage of all surfaces.

- Add a non-ionic surfactant such as Agral or crop oil such as Hasten at recommended rates.
- Target eggs, newly-hatched larvae, nymphs and immature insect life stages. To maximise efficacy, apply sequential treatment at 28 day spray interval.
- DO NOT apply more than 2 applications per crop.
- DO NOT apply less than 28 days after the initial treatment.
- Follow the current Insecticide Resistance Management Strategies for Group 28 insecticides. Refer to resistance management strategies for Silverleaf whitefly and Melon aphid and Green peach aphid.
- Caution must be exercised when applying Minecto Forte with Hasten on sensitive crops during periods when plants are subjected to drought stress, high temperatures or high humidity.
- Minecto Forte is not compatible with IPM.

Withholding Period:

DO NOT use on plants grown for human or animal consumption.

Safety directions:

Harmful if inhaled. May irritate the eyes. Avoid contact with eyes. Do not inhale vapour or spray mist. When opening the container, preparing spray, and using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves, and disposable mist facemask covering mouth and nose. In addition, if applying by airblast or vertical sprayer wear a washable hat. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

First aid instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Re-entry statement:

DO NOT enter treated areas until the spray has dried or install or maintain handset irrigation in treated nursery stock for 10 days after treatment, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Jurisdiction:

All States and Territories.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the Agricultural and Veterinary Chemicals Code Act 1994, in accordance with the obligation imposed by that section.

To Avoid Crop Damage

Minecto Forte has been used on a number of field crops without damage. However, some plant species and varieties are particularly sensitive to chemical products and/or when applied during certain environmental conditions. It is advisable to treat only a small number of plants first, in order to determine their reaction to the product, before large scale use.

Issued by the Australian Pesticides and Veterinary Medicines Authority