



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW EMERGENCY USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF POWDERY MILDEW IN WHEAT**

PERMIT NUMBER – PER93216

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 16 FEBRUARY 2023 TO 31 JULY 2024

Permit Holder:

GRAIN PRODUCERS AUSTRALIA LIMITED
12 Cooma Road
BRAIDWOOD NSW 2622

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

TALENDO FUNGICIDE (APVMA No. 64165)

Containing: 200 g/L PROQUINAZID as the only active constituent.

RESTRAINT:

DO NOT apply with aircraft.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff from the treatment area for at least 3 days after application.

Comply with all spray drift restraints on the product label.

Directions for Use:

Crop	Disease	Rate
Wheat	Powdery Mildew (<i>Blumeria graminis</i> f. <i>Sp tritici</i>)	250 mL/ha

Critical Use Comments:

- Monitor crops regularly from early tillering. Apply early at the first signs of disease (after BBCH 25 - 5 tillers present) as a protectant treatment only.
- Monitor if conditions favour disease development and reapply from 14 to 21 days after the first application.
- Apply foliar application by ground boom.
- Use a spray volume of 100-200 L/ha.
- DO NOT apply more than 2 applications per crop.
- DO NOT apply less than 14 days after the initial treatment.
- DO NOT apply after growth stage BBCH49 (first awns visible – before beginning of heading).
- Talendo fungicide has no curative activity and does not control existing infections (including powdery mildew between infection and visible stage).
- Apply in tank mixture with another approved fungicide with a different Mode of Action (MoA).
- Apply proquinazid in accordance with the current CropLife Fungicide Resistance Management Strategy.

Withholding Period:

Harvest: Not required when used as directed.

Grazing: Do not graze or cut for stock food for 4 weeks after application.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

The sensitivity of the crop to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.